

Accord logo

Package leaflet: Information for the patient

## **Glyceryl Trinitrate tablets 500micrograms and 600micrograms**

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet**

- 1. What Glyceryl Trinitrate tablets are and what they are used for**
- 2. What you need to know before you take Glyceryl Trinitrate tablets**
- 3. How to take Glyceryl Trinitrate Tablets**
- 4. Possible side effects**
- 5. How to store Glyceryl Trinitrate Tablets**
- 6. Contents of the pack and other information**

### **1. What Glyceryl Trinitrate tablets are and what they are used for**

Glyceryl Trinitrate belongs to a group of medicines called nitrate vasodilators. These medicines work by relaxing the blood vessels of the heart. This reduces the strain on the heart by making it easier to pump blood.

Glyceryl Trinitrate tablets may be used to:

- prevent or stop chest pain caused by angina pectoris
- stop spasms in the blood vessels of the heart.

### **2. What you need to know before you take Glyceryl Trinitrate tablets**

**Do not take** Glyceryl Trinitrate tablets if you:

- are **allergic** to glyceryl trinitrate, nitrates or any of the ingredients of this medicine (listed in section 6). An allergic reaction may include a rash, itching, difficulty breathing or swelling of the face, lips, throat or tongue.
- have **anaemia**.
- have **raised pressure in your head** including that caused by an accident to your head or bleeding on the brain, causing painful eyes, changes in vision and a bad headache especially behind the eyes.
- have increased pressure in the eye (**glaucoma**).
- are taking sildenafil (Viagra) or similar medicines (such as tadalafil or vardenafil) used for the **treatment of erection problems**.
- suffer from a heart condition called **hypertrophic obstructive cardiomyopathy**.

### **Warnings and precautions**

**Talk to your doctor or pharmacist** before taking Glyceryl Trinitrate tablets if you have:

- severe **liver** or **kidney** problems
- an **under active thyroid** gland
- had a recent **heart attack**
- **low body temperature** (hypothermia)
- poor nutrition due to an unbalanced diet or digestion problems (**malnutrition**)
- abnormally low levels of oxygen in the blood (**hypoxaemia**)
- very **low blood pressure**, **low blood volume** or conditions that **lower blood volume in the heart**

- any of the following **heart conditions**: narrowing of heart valves, inflammation of the heart, build up of fluid around the heart or heart failure, disease of the blood vessels supplying the brain (**cerebrovascular disease**).

### **Children and adolescents**

Do not give this medicine to children.

### **Other medicines and Glyceryl Trinitrate Tablets**

**Tell your doctor or pharmacist** if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription. Especially:

- **sildenafil** (Viagra), **tadalafil** (Cialis) or **vardeafil** (Levitra) used to treat erection problems
- **heparin** used to thin the blood
- **n-acetylcysteine** used in paracetamol overdose
- **long acting nitrates** (isosorbide dinitrate, isosorbide mononitrate)
- medicines used to **treat high blood pressure**
- medicines that can **cause dry mouth** (amitriptyline, disopyramide, atropine and propantheline, tricyclic antidepressants)
- **apomorphine** used to treat Parkinson's disease.
- medicines used to treat migraine (**ergot alkaloids**).

### **Glyceryl Trinitrate tablets with alcohol**

You are advised **not** to drink alcohol with Glyceryl Trinitrate tablets, as some of the effects of this medicine are increased.

### **Pregnancy and breast feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

### **Driving and using machines**

As glyceryl trinitrate can cause dizziness, you should make sure you are not affected before driving or operating machinery, especially if taken with alcohol.

### **Glyceryl Trinitrate Tablets contain lactose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

## **3. How to take Glyceryl Trinitrate Tablets**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

**Do not swallow the tablets.**

### **The recommended dose is:**

**Adult:** Take the tablets sat down when taking to stop angina pectoris. 1 tablet should be placed under the tongue and allowed to dissolve slowly. This dose should be repeated after 5 minutes if symptoms are not relieved. If pain persists after a total of 3 doses in 15 minutes, you should seek medical advice immediately.

**Older people:** When taking to prevent or stop angina pectoris, take the tablets sat down to reduce possible fainting or low blood pressure.

Your doctor will change the dose to suit your condition. If you notice that your tablets are not working as well at relieving pain as they have done, contact your pharmacist or doctor. If you need more tablets than before to help, tell the doctor who will advise what to do.

### **If you take more Glyceryl Trinitrate Tablets than you should**

If you (or someone else) swallow a lot of tablets at the same time, or you think a child may have swallowed any, contact your nearest hospital casualty department or tell your doctor immediately. Symptoms of an overdose include being sick, restlessness, low blood pressure, fainting, blueness of the skin, methaemoglobin in the blood, breathing difficulties, lack of oxygen in tissues, slow heart beat and mental illness.

#### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**If any of the following serious side effects happen, stop taking the medicine and tell your doctor immediately or go to your nearest hospital:**

Worsening of angina symptoms, chest pains, hypoxaemia (difficulty breathing, increased rate of breathing), methaemoglobinaemia (breathing problems, headache, tiredness), difficulty breathing, blue colouring of the lips, fingers or toes (cyanosis), rapid (tachycardia) or slowed (bradycardia) heartbeat, blindness in one eye, cerebral ischaemia which may cause weakness in one arm or leg, or weakness in one entire side of the body, difficulty speaking, slurred speech, loss of coordination, loss of consciousness, allergic skin reactions.

**Other side effects:**

**Very common (may affect more than 1 in 10 people)**

- throbbing headache.

**Common (may affect up to 1 in 10 people)**

- spinning sensation, fall in blood pressure when seated or standing up, dizziness, drowsiness
- lack of energy.

**Uncommon (may affect up to 1 in 100 people)**

- fainting
- facial flushing
- feeling or being sick
- localised burning sensation
- tongue blistering
- circulatory collapse.

**Very rare (may affect up to 1 in 10,000 people)**

- restlessness
- heartburn
- bad breath, itchy, flaky or red skin, skin rash.

**Not known (frequency cannot be estimated from the available data)**

- pain around the eye, changes in vision
- palpitations
- weakness.

#### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

#### **5. How to store Glyceryl Trinitrate Tablets**

Keep this medicine out of the sight and reach of children.

The active medicine quickly evaporates from the tablet and you should observe the following conditions to ensure continued potency of the tablets.

- If you do not use the tablets within 8 weeks of first opening the bottle, obtain a fresh supply and return the old tablets to your pharmacist. (Fresh tablets should produce a slight burning sensation

when placed under the tongue; if this does not occur, obtain a fresh supply). In any event do not use the tablets after the expiry date shown.

- Your Glyceryl Trinitrate tablets will be in a glass airtight container with a foil-lined cap. Keep them in this.
- After each use, close the bottle cap tightly.
- Do not put cotton wool, other medicines, or anything else in the bottle with the tablets.
- Store the tablets below 25°C in a dry place, protected from light. If you need to carry them with you, **do not** carry them close to your body, rather in a purse, wallet or handbag.

Do not use this medicine after the expiry date stated on the bottle. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Glyceryl Trinitrate tablets contain**

- The active substance (the ingredient that makes the tablets work) is glyceryl trinitrate BP. Each tablet contains either 0.525mg or 0.630mg of the active ingredient Glyceryl trinitrate 2% adsorbate which is equivalent to 500micrograms and 600micrograms of Glyceryl trinitate.
- The other ingredients are lactose, stearic acid, magnesium stearate, colloidal silicon dioxide, mannitol and acacia.

### **What Glyceryl Trinitrate tablets look like and contents of the pack**

Glyceryl Trinitrate tablets 500 micrograms are white, circular, biconvex uncoated tablets, impressed “C” on one face and the identifying letters “GF” on the reverse.

Glyceryl Trinitrate tablets 600 micrograms are white, circular, biconvex uncoated tablets, impressed “C” on one face and the identifying letters “GS” on reverse.

Pack size is 100.

### **Marketing Authorisation Holder and Manufacturer**

Accord, Barnstaple, EX32 8NS, UK

This leaflet was last revised in October 2020

---